

# PubMed

[U.S. National Library of Medicine](#)  
[National Institutes of Health](#)

[J Am Coll Surg](#). 2010 Apr;210(4):480-90.

## **Early healing of transcolonic and transgastric natural orifice transluminal endoscopic surgery access sites.**

[Mathews JC](#), [Chin MS](#), [Fernandez-Esparrach G](#), [Shaikh SN](#), [Pietramaggiori G](#), [Scherer SS](#), [Ryan MB](#), [Ferrigno M](#), [Orgill DP](#), [Thompson CC](#).

Division of Plastic Surgery, Department of Surgery, Brigham and Women's Hospital, Boston, MA 02115, USA.

### **Abstract**

**BACKGROUND:** Natural Orifice Transluminal Endoscopic Surgery (NOTES) is a developing, minimally invasive surgical approach whose potential benefits are being investigated. Little is known about secure access site closure and early healing kinetics of transvisceral access. **STUDY DESIGN:** Transvisceral access incisions were created in the colon (C-NOTES, n = 8) and stomach (G-NOTES, n = 8) for peritoneal exploration. Incisions were closed primarily with endoloops, endoclips, or t-tags. Macroscopic and histologic analyses performed on postoperative day 7 assessed gross appearance, granulation tissue, inflammation, ulceration, and complications. **RESULTS:** Macroscopically, incisions appeared closed without intraperitoneal spillage. Incisions closed by endoloop and t-tags showed intense granulation tissue fill of defect despite partial (G-NOTES, n = 3) and transmural ulceration (C-NOTES, n = 8; G-NOTES, n = 3). Of the 30 t-tags applied, 40% broke or deployed into the peritoneal cavity. Endoclip closures (C-NOTES, n = 1; G-NOTES, n = 1) did not show histologic mucosal continuity. Healing complications included transmural necrosis (C-NOTES, n = 1; G-NOTES, n = 1), foreign body material (C-NOTES, n = 3; G-NOTES, n = 2), and microabscesses (G-NOTES, n = 1). **CONCLUSIONS:** This study provides a reproducible model to assess noninvasive repair of planned visceral perforations. Of investigated technologies, endoloop closure was favored for transcolonic incisions, and t-tags with omental patch for transgastric incisions, although these have significant limitations. Endoclips were inadequate for primary closure, but may be useful as an adjunctive closure modality. Additional studies are needed to examine visceral repair at later time points, as they will help determine the quality and kinetics of repair of a variety of incision closure strategies. This study demonstrates the need for improved technologies to more reliably close

visceral transluminal defects. Copyright (c) 2010 American College of Surgeons. Published by Elsevier Inc. All rights reserved.

PMID: 20347741 [PubMed - indexed for MEDLINE]

## Supplemental Content



### Related articles

- [Transluminal endosurgery: novel use of endoscopic tacks for the closure of access sites in natural orifice transluminal endoscopic surgery \(with videos\).](#)  
*Gastrointest Endosc.* 2009 May; 69(6):1161-6.  
[Gastrointest Endosc. 2009]
- [Evaluation of a novel access and closure device for NOTES applications: a transcolonic survival study in the porcine model \(with video\).](#)  
*Gastrointest Endosc.* 2008 May; 67(6):964-9.  
[Gastrointest Endosc. 2008]
- [Air and fluid leak tests after NOTES procedures: a pilot study in a live porcine model \(with videos\).](#)  
*Gastrointest Endosc.* 2008 Sep; 68(3):513-9. Epub 2008 Apr 9.  
[Gastrointest Endosc. 2008]
- [Endoscopic closure of the natural orifice transluminal endoscopic surgery \(NOTES\) access site to the peritoneal cavity by means of transmural resorbable sutures: an animal survival study.](#)  
*Endoscopy.* 2009 Feb; 41(2):154-9. Epub 2009 Feb 12.  
[Endoscopy. 2009]
- [Review A primer on natural orifice transluminal endoscopic surgery: building a new paradigm.](#)  
*Surg Innov.* 2006 Jun; 13(2):86-93.  
[Surg Innov. 2006]
- » [See reviews...](#) | » [See all...](#)